

Math. J. Okayama Univ. **63** (2021), 53–60

REMARK ON A PAPER BY IZADI AND BAGHALAGHDAM ABOUT CUBES AND FIFTH POWERS SUMS

GAKU IOKIBE

ABSTRACT. In this paper, we refine the method introduced by Izadi and Baghalaghdam to search integer solutions to the Diophantine equation $X_1^5 + X_2^5 + X_3^5 = Y_1^3 + Y_2^3 + Y_3^3$. We show that the Diophantine equation has infinitely many positive solutions.

1. INTRODUCTION

In [2], Izadi and Baghalaghdam consider the Diophantine equation:

$$(1) \quad a(X_1'^5 + X_2'^5) + \sum_{i=0}^n a_i X_i^5 = b(Y_1'^3 + Y_2'^3) + \sum_{i=0}^m b_i Y_i^3$$

where $n, m \in \mathbb{N} \cup \{0\}$, $a, b \neq 0$, a_i, b_i are fixed arbitrary rational numbers. They use theory of elliptic curves to find nontrivial integer solutions to (1). In particular, they discuss the equation:

$$(2) \quad X_1^5 + X_2^5 + X_3^5 = Y_1^3 + Y_2^3 + Y_3^3$$

and obtain integer solutions, for example:

$$8^5 + 6^5 + 14^5 = (-110)^3 + 124^3 + 14^3,$$

$$128122^5 + (-79524)^5 + 48598^5 = 359227580^3 + (-251874598)^3 + 107352982^3.$$

However, no positive solutions are presented in their paper [2]. In this paper, we refine their method to find positive solutions to (2).

Consider the Diophantine equation (2). Let:

$$(3) \quad \begin{cases} X_1 = t + x_1, & X_2 = t - x_1, & X_3 = \alpha t, \\ Y_1 = t + v, & Y_2 = t - v, & Y_3 = \beta t. \end{cases}$$

Then we get a quartic curve:

$$(4) \quad C : v^2 = \frac{2 + \alpha^5}{6} t^4 + \frac{20x_1^2 - 2 - \beta^3}{6} t^2 + \frac{5x_1^4}{3}$$

with parameters $x_1, \alpha, \beta \in \mathbb{Q}$. If we get a rational point (t, v) on C , we can compute a rational solution to (2) (see [2]).

Mathematics Subject Classification. Primary 11D41; Secondary 11D45, 14H52.

Key words and phrases. Diophantine equations, Elliptic Curves.

Once we obtain rational solutions to (2), we can obtain integer solutions by multiplying an appropriate value to X_i, Y_i . In the same way, in order to obtain solutions in positive integers, it suffices to search positive rational solutions to equation (2).

2. ADDITIONAL REQUIREMENTS FOR POSITIVE SOLUTIONS

Suppose that a positive rational solution $(X_i, Y_i)_{1 \leq i \leq 3}$ to (2) is obtained from a given point (t, v) on the quartic C .

Proposition 2.1. Let $\alpha, \beta, x_1 \in \mathbb{Q}$ and

$$F(t) = \frac{2 + \alpha^5}{6}t^4 + \frac{20x_1^2 - 2 - \beta^3}{6}t^2 + \frac{5x_1^4}{3}.$$

A rational point (t, v) on the curve $C : v^2 = F(t)$ in (4) produces a positive rational solution to (2) by (3) if and only if

$$(5) \quad \alpha, \beta > 0, \quad 0 \leq F(t) < t^2, \quad t > |x_1|$$

hold.

Proof. If X_i and Y_i are positive in the solution in the form (3), we have $t = (X_1 + X_2)/2 > 0$, $\alpha = 2X_3/(X_1 + X_2) > 0$ and $\beta = 2Y_3/(Y_1 + Y_2) > 0$. For $(t, v) \in C$, one has that $0 \leq v^2 = F(t) < v^2 + Y_1Y_2 = t^2$. It follows from $x_1^2 < x_1^2 + X_1X_2 = t^2$ that $t > |x_1|$ for $t > 0$. Conversely, suppose the inequalities in (5) hold. Then the given point (t, v) on C satisfies $v^2 = F(t) < t^2$. This and (5) immediately imply $X_i, Y_i > 0$ in (3). \square

Proposition 2.2. Under the same assumption as Proposition 2.1, let

$$(6) \quad a = \frac{2 + \alpha^5}{6}, \quad b = \frac{20x_1^2 - 8 - \beta^3}{6}, \quad c = \frac{5}{3}x_1^4.$$

Then a, b, c satisfy $b^2 - 4ac > 0$ and $b < 0$ if and only if there exists a real number t such that $F(t) < t^2$.

Proof. Let $\tilde{F}(t) = F(t) - t^2$. Since $\tilde{F}(0) = 5x_1^4/3 \geq 0$, and in this case $a > 0$, it is easy to see that the following conditions are equivalent to each others:

- (i) There exists a real number t such that $F(t) < t^2$.
- (ii) The equation $\tilde{F}(t) = 0$ has four distinct solutions.
- (iii) The quadratic equation $ax^2 + bx + c = 0$ has two distinct non-negative solutions.
- (iv) The discriminant $D = b^2 - 4ac$ of the quadratic function $f(x) = ax^2 + bx + c$ is positive, and the axis of the quadratic function $-b/2a$ is positive, and $f(0) \geq 0$.

The condition (iv) holds if and only if “ $b^2 - 4ac > 0$ and $b < 0$ ”, since $a > 0$ and $f(0) = c = 5x_1^4/3 \geq 0$. \square

3. EXAMPLE FOR $X_1^5 + X_2^5 + X_3^5 = Y_1^3 + Y_2^3 + Y_3^3$

Let us first search parameters (x_1, α, β) such that

$$0 < \alpha, \beta, \quad b < 0 < b^2 - 4ac$$

with a, b, c given by (6) and such that the quartic curve C of (4) has at least one rational point. Note that these are necessary to satisfy conditions of Proposition 2.1, 2.2. Then, the curve C is birationally equivalent to an elliptic curve E over \mathbb{Q} . If E has positive rank, then C has infinitely many rational points.

Let $(x_1, \alpha, \beta) = (2, 1, 16)$. Then the quartic:

$$C : v^2 = \frac{1}{2}t^4 - \frac{2009}{3}t^2 + \frac{80}{3},$$

has a rational point $(t, v) = (44, 760)$. By $T = t - 44$, we transform C into

$$C' : v^2 = \frac{1}{2}T^4 + 88T^3 + \frac{15415}{3}T^2 + \frac{334312}{3}T + 760^2$$

which is birationally equivalent over \mathbb{Q} to the cubic elliptic curve (see [5, Theorem 2.17], [2]):

$$E : y^2 + \frac{41789}{285}xy + 133760y = x^3 - \frac{76876021}{324900}x^2 - 1155200x + \frac{2460032672}{9},$$

where:

$$T = \frac{2 \cdot 760(x + \frac{15415}{3}) - \frac{334312^2}{2 \cdot 3^2 \cdot 760}}{y}, \quad v = -760 + \frac{T(Tx - \frac{334312}{3})}{2 \cdot 760}.$$

Using the Sage software [3], we find that the cubic curve E is an elliptic curve which has rank 2 and the generators of E are:

$$P_1 = \left(-\frac{1802189}{1521}, \frac{5513659679}{417430} \right), \quad P_2 = \left(-\frac{351379}{363}, \frac{47356344241}{2276010} \right).$$

We now consider the subset

$$C_0 = \{(t, v) \in C \mid 0 \leq F(t) < t^2\} \subset C$$

whose points satisfy another condition (5) of Proposition 2.1. The two quartic equations:

$$F(t) = \frac{1}{2}t^4 - \frac{2009}{3}t^2 + \frac{80}{3} = 0, \quad \tilde{F}(t) = \frac{1}{2}t^4 - \frac{2009}{3}t^2 + \frac{80}{3} - t^2 = 0$$

have respectively solutions:

$$t = \pm \frac{1}{3} \sqrt{6027 \pm 3\sqrt{4035601}}, \quad t = \pm \frac{2}{3} \sqrt{1509 \pm 3\sqrt{252979}}.$$

Let us take larger solutions as:

$$\begin{aligned} a_1 &= \frac{1}{3}\sqrt{6027 + 3\sqrt{4035601}} \simeq 36.59635926... \\ a_2 &= \frac{2}{3}\sqrt{1509 + 3\sqrt{252979}} \simeq 36.62367500... \end{aligned}$$

If a point (t_0, v_0) on C satisfies $a_1 \leq t_0 \leq a_2$, then (t_0, v_0) lies on C_0 . We now make use of the composition law of points on the elliptic curve E . Since E has positive rank, we can test infinitely many rational points of E till finding a point (t_0, v_0) on C_0 . We find that the rational point

$$Q = 2P_1 - P_2 = \left(\frac{304845381192111829037}{58470412871306667}, -\frac{4767546475726965161322288395890039}{4652843756178203561643745770} \right)$$

on E corresponds to

$$(t_0, v_0) = \left(\frac{170815619844155909156204}{4664941095250009917983}, -\frac{690740884062625663919872925291699877683029096}{21761675422152362106175457381859866386788289} \right)$$

on C_0 , and creates a positive rational solution:

$$\begin{aligned} X_1 &= \frac{180145502034655928992170}{4664941095250009917983} \simeq 38.61688676... \\ X_2 &= \frac{161485737653655889320238}{4664941095250009917983} \simeq 34.61688676... \\ X_3 &= \frac{170815619844155909156204}{4664941095250009917983} \simeq 36.61688676... \\ Y_1 &= \frac{106103920658980331397442614601687483092587436}{21761675422152362106175457381859866386788289} \simeq 4.875723886... \\ Y_2 &= \frac{1487585688784231659237188465185087238458645628}{21761675422152362106175457381859866386788289} \simeq 68.35804964... \\ Y_3 &= \frac{2733049917506494546499264}{4664941095250009917983} \simeq 585.8701882... \end{aligned}$$

Next we shall prove that the Diophantine equation (2) has infinitely many positive solutions. The real locus of elliptic curve $E(\mathbb{R})$ can be regarded as a compact topological subspace of complex projective variety E .

Lemma 3.1. If the rank of elliptic curve E over \mathbb{Q} is positive, every point of $E(\mathbb{Q})$ is an accumulation point in $E(\mathbb{R})$.

Proof. Since $E(\mathbb{R})$ is a compact topological group, and $E(\mathbb{Q})$ is an infinite subgroup of $E(\mathbb{R})$, there is at least one accumulation point of $E(\mathbb{Q})$ in $E(\mathbb{R})$. The group operations are homeomorphisms from $E(\mathbb{R})$ to itself. Therefore all points of $E(\mathbb{Q})$ are accumulation points of $E(\mathbb{R})$. \square

Theorem 3.2. The Diophantine equation (2) has infinitely many positive solutions.

Proof. The part of C_0 has one rational point (t_0, v_0) which corresponds to the above point Q . By Lemma 3.1, the point Q is an accumulation point of $E(\mathbb{Q})$ in $E(\mathbb{R})$, and (t_0, v_0) is that of $C(\mathbb{Q})$ in $C(\mathbb{R})$. Thus the part of C_0 includes infinitely many rational points. Since $2 = |x_1| < a_1 = 36.59635926...$, they correspond to positive rational solutions to (2). \square

4. EXAMPLE FOR $X_1^5 + X_2^5 = Y_1^3 + Y_2^3 + Y_3^3$

Let $\alpha = 0$. Then (2) gives another Diophantine equation:

$$(7) \quad X_1^5 + X_2^5 = Y_1^3 + Y_2^3 + Y_3^3.$$

In the same way, we can obtain a rational or positive rational solutions of it. For example, let $x_1 = 10$, $\beta = 18$. Then the quartic curve:

$$C : v^2 = \frac{1}{3}t^4 - 639t^2 + \frac{50000}{3}$$

has a rational point $(t, v) = (-5, 30)$ and can be regarded as an elliptic curve over \mathbb{Q} that has rank 2. It is birationally equivalent to:

$$E : y^2 + \frac{1867}{9}xy - 400y = x^3 - \frac{3676525}{324}x^2 - 1200x + \frac{367652500}{27}.$$

From this, we can compute positive rational solutions to (7). For example, there is a point $Q = (x_0, y_0)$ on E with

$$x_0 = \frac{9233921838917810856046138588468998730}{71226852166762122405616706766475947}$$

corresponding to (t_0, v_0) on C with

$$t_0 = \frac{7869911761727476320751662986237524106650}{180965667579279848488380712753242417827}$$

which creates the following solution to (7):

$$\begin{aligned} X_1 &= \frac{9679568437520274805635470113769948284920}{180965667579279848488380712753242417827}, \\ X_2 &= \frac{60602550859346778358678558705099928380}{180965667579279848488380712753242417827}, \\ Y_1 &= \frac{2102579397586077496858869804126511993988094601307100986270258503567645177035000}{32748572842414417658282657731373155447687070419319181813277645661864847401929}, \\ Y_2 &= \frac{745788273916000738265027095213285105579870143595196644344754733646843241464100}{32748572842414417658282657731373155447687070419319181813277645661864847401929}, \\ Y_3 &= \frac{141658411711094573773529933752275433919700}{180965667579279848488380712753242417827}. \end{aligned}$$

The case of $\beta = 0$ will be discussed briefly in 5.2 below.

5. PARAMETERS (x_1, α, β) FROM TRIVIAL SOLUTIONS

5.1. There are several trivial solutions; for example:

$$1^5 + 1^5 + 1^5 = 1^3 + 1^3 + 1^3.$$

We call solutions to (2) which consist of 0, ± 1 trivial. We are going to check some of them to search integer (or positive) solutions.

A solution to (2) may decide parameter. For example, when $X_i = Y_i = 1$ ($i = 1, 2, 3$), we get $(x_1, \alpha, \beta) = (0, 1, 1)$. Then:

$$C : v^2 = \frac{1}{2}t^4 - \frac{1}{2}t^2$$

has a singular point $(t, v) = (0, 0)$ and can be parametrized by one parameter. Let us divide both sides of C by t^4 and substitute s, w for $1/t, v/t^2$ respectively. Then:

$$C' : w^2 = \frac{1}{2} - \frac{1}{2}s^2$$

has a rational point $(s, w) = (1, 0)$. Hence we can parametrize rational points on C' and integer solutions to (2). That is to say we have:

$$\begin{aligned} & \left(\frac{2k^2 + 1}{2k^2 - 1} \right)^5 + \left(\frac{2k^2 + 1}{2k^2 - 1} \right)^5 + \left(\frac{2k^2 + 1}{2k^2 - 1} \right)^5 \\ &= \left(\frac{4k^4 - 4k^3 - 2k - 1}{(2k^2 - 1)^2} \right)^3 + \left(\frac{4k^4 + 4k^3 + 2k - 1}{(2k^2 - 1)^2} \right)^3 + \left(\frac{2k^2 + 1}{2k^2 - 1} \right)^3 \end{aligned}$$

where $k \in \mathbb{Q}$. We can see that large enough k give positive solutions to (2). For example:

$$\left(\frac{9}{7} \right)^5 + \left(\frac{9}{7} \right)^5 + \left(\frac{9}{7} \right)^5 = \left(\frac{27}{49} \right)^3 + \left(\frac{99}{49} \right)^3 + \left(\frac{9}{7} \right)^3$$

where $k = 2$. Since $X_1 = X_2 = X_3 = Y_3$, this solution also gives positive solution to another Diophantine equation $3X^5 = Y_1^3 + Y_2^3 + X^3$. Moreover it satisfies $X_1 + X_2 + X_3 = Y_1 + Y_2 + Y_3$ because $\alpha = \beta$.

5.2. From another trivial solution:

$$1^5 + 0^5 + 0^5 = 1^3 + 0^3 + 0^3,$$

we can derive parameters $(x_1, \alpha, \beta) = (\frac{1}{2}, 0, 0)$. Then:

$$C : v^2 = \frac{1}{3}t^4 + \frac{1}{2}t^2 + \frac{5}{48}$$

is an elliptic curve defined over \mathbb{Q} with rational point $(t, v) = (\frac{1}{2}, \frac{1}{2})$. It is birationally equivalent to:

$$E : y^2 + \frac{4}{3}xy + \frac{2}{3}y = x^3 + \frac{5}{9}x^2 - \frac{1}{3}x - \frac{5}{27}$$

over \mathbb{Q} and has rank 1. Hence we can apply the method of Section 3 to compute positive solutions to

$$(8) \quad X_1^5 + X_2^5 = Y_1^3 + Y_2^3$$

as a special case of (2) with $X_1, X_2, Y_1, Y_2 > 0, X_3 = Y_3 = 0$ (where $\alpha = \beta = 0$ in (3)). For example, a point

$$Q = \left(\frac{10017045137918654785}{165672066306928896}, \frac{29224609136538294659462738431}{67433225470590933809197056} \right)$$

on E corresponding to the point

$$(t_0, v_0) = \left(\frac{2806052350871126431439}{4379016004568066987998}, \frac{5797926783162005502807971914786692611082209}{9587890584131638439948667971418559938024002} \right)$$

on C creates the positive solution to (8):

$$\begin{aligned} X_1 &= \frac{2497780176577579962719}{2189508002284033493999}, \quad X_2 = \frac{308272174293546468720}{2189508002284033493999}, \\ Y_1 &= \frac{5970900430111130674379700360675596051258385}{4793945292065819219974333985709279969012001}, \\ Y_2 &= \frac{172973646949125171571728445888903440176176}{4793945292065819219974333985709279969012001}. \end{aligned}$$

5.3. There exists one more parameter with $\beta = 0$, $(x_1, \alpha, \beta) = (0, 0, 0)$, which is derived from the trivial solution:

$$1^5 + 1^5 + 0^5 = 1^3 + 1^3 + 0^3.$$

Then the rational points on:

$$C : v^2 = \frac{1}{3}t^4 - \frac{1}{3}t^2$$

can be parametrized. Thus we have:

$$\left(\frac{3k^2+1}{3k^2-1}\right)^5 + \left(\frac{3k^2+1}{3k^2-1}\right)^5 = \left(\frac{9k^4-6k^3-2k-1}{(3k^2-1)^2}\right)^3 + \left(\frac{9k^4+6k^3+2k-1}{(3k^2-1)^2}\right)^3,$$

where $k \in \mathbb{Q}$. For example, substituting 2 for k , we have:

$$\left(\frac{13}{11}\right)^5 + \left(\frac{13}{11}\right)^5 + 0^5 = \left(\frac{91}{121}\right)^3 + \left(\frac{195}{121}\right)^3 + 0^3.$$

The solutions which are obtained in these ways give solutions to another Diophantine equation $2X^5 = Y_1^3 + Y_2^3$.

5.4. It is not simple to find parameters (x_1, α, β) that produce elliptic curves for non-trivial solutions $(X_i, Y_i)_{1 \leq i \leq 3}$. In particular, the author could not find a good parameter for $\beta = 0$, $\alpha \neq 0$:

Question 5.1. Find (a good method for) positive solutions to:

$$X_1^5 + X_2^5 + X_3^5 = Y_1^3 + Y_2^3.$$

Acknowledgement: The author would like to thank the referee for many valuable suggestions to improve this article.

REFERENCES

- [1] G.Iokibe: *Search for positive solutions to Diophantine equations with cubes and fifth powers sums*, Master thesis, Department of Mathematics, Osaka University, February 2018.
- [2] F.Izadi and M.Baghalaghdam: *On the Diophantine equation in the form that a sum of cubes equals a sum of quintics*, Math. J. Okayama Univ. **61** (2019), 75–84. (arXiv:1704.00600v1 [math.NT] 30 Mar 2017)
- [3] Sage software, available from <http://www.sagemath.org/>
- [4] J. Silverman: *The Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, second edition, 2008.

- [5] L. C. Washington: *Elliptic Curves: Number Theory and Cryptography*, Chapman & Hall/CRC, 2003.

GAKU IOKIBE

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY,
TOYONAKA, OSAKA 560-0043, JAPAN

e-mail address: u325137g@alumni.osaka-u.ac.jp

(Received December 16, 2018)

(Accepted June 19, 2019)